

1. Cheng, Y.-L., Kuan, J.-E., Wang, C.-W., Chen, R.-H. 2025. Mga2-mediated transcription supports mitotic nuclear expansion under lipid saturation conditions in stearoyl-CoA desaturase Ole1 mutant. ***Mol. Biol. Cell*** 36(11):ar144, 1-17.
2. Huang, L.-J. and Chen, R.-H. 2022. Lipid saturation induces degradation of squalene epoxidase for sterol homeostasis and cell survival. ***Life Sci. Alliance*** 6(1):e202201612.
3. Chen, R.-H. 2019. Chromosome detachment from the nuclear envelope is required for genome stability in closed mitosis. ***Mol. Biol. Cell*** 30(13):1578-1586.
4. Hsu, T.-H., Chen, R.-H., Cheng, Y.-H., and Wang, C.-W. 2017. Lipid droplets are central organelles for meiosis II progression during yeast sporulation. ***Mol. Biol. Cell*** 28(3): 440-451.
5. Yang, P.-L., Hsu, T.-H., Wang, C.-W., and Chen, R.-H. 2016. Lipid droplets maintain lipid homeostasis during anaphase for efficient cell separation in budding yeast. ***Mol. Biol. Cell*** 27(15):2368-2380.
6. Cheng, Y.-L. and Chen, R.-H. 2015. Assembly and quality control of the protein phosphatase 1 holoenzyme involves the Cdc48-Shp1 chaperone. ***J. Cell Sci.*** 128(6):1180-1192.
7. Madsen, L., Molbæk, K., Larsen, I.B., Nielsen, S.V., Poulsen, E.G., Walmod, P.S., Hofmann, K., Seeger, M., Chien, C.Y., Chen, R.-H., Kriegenburg, F., Hartmann-Petersen, R. 2014. Human ASPL/TUG interacts with p97 and complements the proteasome mislocalization of a yeast ubx4 mutant, but not the ER-associated degradation defect. ***BMC Cell Biol.*** 15:31.
8. Chien, C.-Y. and Chen, R.-H. 2013. Cdc48 chaperone and adaptor Ubx4 distribute the proteasome in the nucleus for anaphase proteolysis. ***J. Biol. Chem.*** 288(52): 37180-37191.
9. Chen, R.-H. 2012. Nuclear envelope assembly and disassembly during the cell cycle. eLS. John Wiley & Sons, Ltd: Chichester.
10. Tseng, L.-C. and Chen, R.-H. 2011. Temporal control of nuclear envelope assembly by phosphorylation of lamin B receptor. ***Mol. Biol. Cell*** 22:3306-3317.
11. Hsieh, M.-T. and Chen, R.-H. 2011. Cdc48 and cofactors Npl4-Ufd1 are

important for G1 progression during heat stress by maintaining cell wall integrity in *Saccharomyces cerevisiae*. ***PLoS ONE*** 6: e18988.

12. Cheng, Y.-L. and Chen, R.-H. 2010. AAA-ATPase Cdc48 and Cofactor Shp1 promote chromosome bi-orientation by balancing Aurora B activity. ***J. Cell. Sci.*** 123: 2025-2034.